



(a)

FTD-ID(RS)T-0849-87

AD-A187 441

OTIC FILE COPY

FOREIGN TECHNOLOGY DIVISION



INFLUENCE OF COMPLEXING ON THE REACTIVITY OF POLAR UNSATURATED MONOMERS by

B.L. Yerusalimskiy, V.N. Krasulina, Yu. Ye. Eyzner





87 12 11 003

Approved for public release; Distribution unlimited.

PARTIALLY EDITED MACHINE TRANSLATION

FTD-ID(RS)T-0849-87

18 November 1987

MICROFICHE NR: FTD-87-C-001012

INFLUENCE OF COMPLEXING ON THE REACTIVITY OF POLAR UNSATURATED MONOMERS

By: B.L. Yerusalimskiy, V.N. Krasulina, Yu. Ye. Eyzner

English pages: 3

Source: Vysokomolekulyarnyye Soyedineniya, Seriya B,

Kratkiye Soobshcheniya, Vol. 12, Nr. 5,

May 1970, pp. 327-328

Country of origin: USSR

This document is a machine translation.

Input and Merged by: Janet L. Fox

Approved for public release; Distribution unlimited.



Accesi	on For		
סודם	oblined	000	
By			
Availability Codes			
Dist	Aveil and/or Special		
A-1			

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION FOREIGN TECHNOLOGY DIVISION WPAFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Blo	ck	Italic	Transliteration	Block	Italic	Transliteration
Α	a	A a	А, а	Рр	PP	R, r
Б	б	56	B, b	Сс	Cc	S, s
В	В	B •	V, v	Тτ	T m	T, t
Γ	۲	Γ *	G, g	Уу	у у	U, u
Д	щ	Дð	D, d	Фф	Φ φ	F, f
E	е	E e	Ye, ye; E, e*	Х×	X x	Kh, kh
Ж	ж	Ж ж	Zh, zh	Цц	Ll u	Ts, ts
3	э.	3 ,	Z, z	4 4	4 v	Ch, ch
И	И	Н и	I, i	Шш	Ш ш -	Sh, sh
Й	й	A a	Y, y	Щщ	Щ щ	Shch, shch
Н	н	KK	K, k	Ъъ	ъ.	11
Л	л	ЛА	L, 1	Ыы	W ii	Y, y
M	M	Мм	M, m	Ьь	ь.	•
Н	н	Н н	N, n	Ээ	9 ,	Е, е
Ü	0	0 0	0, 0	Юю	10 n	Yu, yu
П	п	Пп	P, p	Яя	Яя	Ya, ya

*ye initially, after vowels, and after b, b; e elsewhere. When written as e in Russian, transliterate as ye or e.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh ⁻¹
cos	cos	ch	cosh	arc ch	cosh_1
tg	tan	th	tanh	arc th	tanh ^l
ctg	cot	cth	coth	arc cth	coth_1
sec	sec	sch	sech	arc sch	sech_1
cosec	CSC	csch	csch	arc csch	csch ⁻¹

Russian	English	
rot	curl	
1 g	log	

GRAPHICS DISCLAIMER

All figures, graphics, tables, equations, etc. merged into this translation were extracted from the best quality copy available.

INFLUENCE OF COMPLEXING ON THE REACTIVITY OF POLAR UNSATURATED MONOMERS

B. L. Yerusalimskiy, V. N. Krasulina, Yu. Ye. Eyzner

Dear editor!

The known data about the behavior of vinyl monomers (M) during anionic polymerization conform to the following order of activity: nitroethylene > acrylonitrile (AN) > methylacrylate (MA) > methyl methacrylate. At the same time the calculations of distribution of electron energy, made by us according to the method of zero differential overlapping [1], lead to another order, as this show & components of charge of carbon atom of vinyl group in the monomers indicated.

$$\frac{+0.033}{\text{CH}_{3}} \times \frac{+0.092}{\text{COOCH}_{2}} \times \frac{+0.060}{\text{CH}_{3}} \times \frac{+0.016}{\text{CH}_{3}} \times \frac{+0.016}{\text{CH}_{$$

Copolymerization of AN with MA in toluene at -60°.
(Molar ratio AN:MA=1. Concentration of lithium butyl 0.01 mole/1).

(//) [ДМБ], мель/л	(2) Выход сопо- яниера, %	ЖА в сопо- жимере, %	
0,01 0,03	7,0 17,6 10,6	36,6 57,3 70,0	

Key: (1). [DMB], mole/l. (2). Yield of copolymer, %. (3). MA
in copolymer, %.

Page 328

contil

We assumed that this nonconformity is caused by different relative contribution of the course of reaction of propagation for each of the monomers through the stage of complexing with the gegenion of the monomers through the stage of complexing with the gegenion of a significant change of the electron density in the viny of group of monomer with the formation of complexes of the type $CH_a = CHX \mapsto MeR$ (where X - polar group, MeR - metal-alkyl) testify our calculations, which relate to the complexes AN and MA with lithiummethyl [2]. The role of complexing as the factor, which influences monomer activity, appeared during the comparison of copolymers - AN-MA, obtained by us under the action of lithiumbutyl and its complex with 2,3-dimethoxybutane (DMB). As it turned out, the coordination saturation of gegenion by the independent base of Lewis leads to the formation of copolymer with the preferred content MA (table), i.e. such composition, which will agree with the results of quantum-chemical calculations.

Apparently, with the formation of complexes $M \to MeR$ is important not only selection of one of the monomers, but also its activation, caused by the change of the electronic characteristic of double bond [2], since the reaction of propagation, which takes place besides the stage of complexing, is not excluded.

Received by the editorial staff 16.I.1970.

Literature.

J. A. Pople, G. A. Segal, J. Chem. Phys., 44, 3289, 1966.
 Б. Л. Ерусалимский, International Symposium on Macromolecular Chemistry, Plenary and Main Lectures, Sect. 3, Budapest, 1970.

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

ORGANIZATION	MICROFICHE
A205 DMAHTC	1
A210 DHAAC	i
BJ44 DIA/RTS-2C	9
CO43 USAMIIA	i
CS00 TRADOC	ī
CSO9 BALLISTIC RES LAB	i
CS10 BAT LABS/AVRADCOM	ī
CS13 ABRADCOM	ī
C535 AVRADCOM/TSARCOM	ī
C539 TRASANA	ī
C591 FSTC	Ž
C619 MIA REDSTONE	1
DOOR WISC	ī
EOSJ MQ USAF/INET	ī
E404 AEDC/DOF	i
E408 AFVIL	i
E410 AD/IND	i
E429 SD/1MD	ī.
POOS DOE/ISA/DDI	ĭ
POSO CIA/OCR/ADD/SD	ž
AFI7/LDE	i
FTD	•
	1
WZA/PHS	i
LLYL/Code L-389	ī
NASA/NST-44	1
XSA/1213/TDL	2
ASD/FTD/YQLA	•
	i i

H M L FED. 1988 DTIC